DDA3020 Machine Learning Lecture 03 Linear Algebra

JIA, Kui School of Data Science, CUHK-SZ

September 14/19, 2023



2 Matrix inverse, determinant, independence



References for this lecture:

- [Book1] Stephen Boyd and Lieven Vandenberghe, "Introduction to Applied Linear Algebra", Cambridge University Press, 2018 (available online).
- [Book2] Andreas C. Muller and Sarah Guido, "Introduction to Machine Learning with Python: A Guide for Data Scientists", O'Reilly Media, Inc., 2017.



2 Matrix inverse, determinant, independence

3 Systems of linear equations

Notations, vectors, matrices

- A scalar is a simple numerical value, like 15 or -3.2.
- Variables or constants that take scalar values are denoted by an italic letter, like x or a.
- We shall focus on real numbers.
- The summation over a collection $\{x_1, x_2, x_3, \ldots, x_m\}$ is denoted like this:

$$\sum_{i=1}^{m} x_i = x_1 + x_2 + \ldots + x_m$$

• The product over a collection $\{x_1, x_2, x_3, \dots, x_m\}$ is denoted like this:

$$\prod_{i=1}^m x_i = x_1 \cdot x_2 \cdot \ldots \cdot x_m$$

- A vector is an ordered list of scalar values, called attributes. We denote a vector as a **bold character**, for example, \mathbf{x} or \mathbf{w} .
- Vectors can be visualized as arrows that point to some directions as well as points in a multi-dimensional space.
- In many books, vectors are written column-wise:

$$\mathbf{a} = \begin{bmatrix} 2\\ 3 \end{bmatrix}, \qquad \mathbf{b} = \begin{bmatrix} -2\\ 5 \end{bmatrix}, \qquad \mathbf{c} = \begin{bmatrix} 1\\ 0 \end{bmatrix}$$

Notations, vectors, matrices

Illustrations of three two-dimensional vectors, $\mathbf{a} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$, $\mathbf{b} = \begin{bmatrix} -2 \\ 5 \end{bmatrix}$, $\mathbf{c} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ are given in Figure 1 below.



Figure: Three vectors visualized as directions and as points.

- We denote an attribute of a vector as an italic value with an index, like this: $w^{(j)}$ or $x^{(j)}$. The index j denotes a specific **dimension** of the vector, the position of an attribute in the list.
- For instance, in the vector **a** shown in red in Figure 1,

$$\mathbf{a} = \begin{bmatrix} a^{(1)} \\ a^{(2)} \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$$
, or more commonly, $\mathbf{a} = \begin{bmatrix} a_1 \\ a_2 \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$

Note:

- The notation $x^{(j)}$ should not be confused with the power operator, such as the 2 in x^2 (squared) or 3 in x^3 (cubed).
- If we want to apply a power operator, say square, to an indexed attribute of a vector, we write like this: $(x^{(j)})^2$.

• A matrix is a rectangular array of numbers arranged in rows and columns. Below is an example of a matrix with two rows and three columns,

$$\mathbf{X} = \begin{bmatrix} 2 & 4 & -3\\ 21 & -6 & -1 \end{bmatrix}$$

• Matrices are denoted with bold capital letters, such as **X** or **W**. **Note:**

- A variable can have two or more indices, like this: $x_i^{(j)}$ or like this $x_{i,j}^{(k)}$.
- For example, in neural networks, we denote as $x_{l,u}^{(j)}$ the input feature j of unit u in layer l.

Operations on Vectors:

$$\mathbf{x} + \mathbf{y} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} x_1 + y_1 \\ x_2 + y_2 \end{bmatrix}$$
$$\mathbf{x} - \mathbf{y} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} - \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} x_1 - y_1 \\ x_2 - y_2 \end{bmatrix}$$
$$a\mathbf{x} = a \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} ax_1 \\ ax_2 \end{bmatrix}$$
$$\frac{1}{a}\mathbf{x} = \frac{1}{a} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} \frac{1}{a}x_1 \\ \frac{1}{a}x_2 \end{bmatrix}$$

• Matrix or Vector Transpose:

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}, \quad \mathbf{x}^{\top} = \begin{bmatrix} x_1 & x_2 \end{bmatrix}$$
$$\mathbf{X} = \begin{bmatrix} x_{1,1} & x_{1,2} & x_{1,3} \\ x_{2,1} & x_{2,2} & x_{2,3} \\ x_{3,1} & x_{3,2} & x_{3,3} \end{bmatrix}, \quad \mathbf{X}^{\top} = \begin{bmatrix} x_{1,1} & x_{2,1} & x_{3,1} \\ x_{1,2} & x_{2,2} & x_{3,2} \\ x_{1,3} & x_{2,3} & x_{3,3} \end{bmatrix}$$

• Dot Product or Inner Product of Vectors:

$$\mathbf{x} \cdot \mathbf{y} = \mathbf{x}^{\top} \mathbf{y}$$
$$= \begin{bmatrix} x_1 & x_2 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}$$
$$= x_1 y_1 + x_2 y_2$$

• Matrix-Vector Product

$$\mathbf{Xw} = \begin{bmatrix} x_{1,1} & x_{1,2} & x_{1,3} \\ x_{2,1} & x_{2,2} & x_{2,3} \\ x_{3,1} & x_{3,2} & x_{3,3} \end{bmatrix} \begin{bmatrix} w_1 \\ w_2 \\ w_3 \end{bmatrix}$$
$$= \begin{bmatrix} x_{1,1}w_1 + x_{1,2}w_2 + x_{1,3}w_3 \\ x_{2,1}w_1 + x_{2,2}w_2 + x_{2,3}w_3 \\ x_{3,1}w_1 + x_{3,2}w_2 + x_{3,3}w_3 \end{bmatrix}$$

Matrix-Vector Product

$$\mathbf{x}^{\top}\mathbf{W} = \begin{bmatrix} x_1 & x_2 & x_3 \end{bmatrix} \begin{bmatrix} w_{1,1} & w_{1,2} & w_{1,3} \\ w_{2,1} & w_{2,2} & w_{2,3} \\ w_{3,1} & w_{3,2} & w_{3,3} \end{bmatrix}$$
$$= \begin{bmatrix} (w_{1,1}x_1 + w_{2,1}x_2 + w_{3,1}x_3) & (w_{1,2}x_1 + w_{2,2}x_2 + w_{3,2}x_3) & (w_{1,3}x_1 + w_{2,3}x_2 + w_{3,3}x_3) \end{bmatrix}$$

Matrix-Matrix Product

$$\mathbf{XW} = \begin{bmatrix} x_{1,1} & \dots & x_{1,d} \\ \vdots & \ddots & \vdots \\ x_{m,1} & \dots & x_{m,d} \end{bmatrix} \begin{bmatrix} w_{1,1} & \dots & w_{1,h} \\ \vdots & \ddots & \vdots \\ w_{d,1} & \dots & w_{d,h} \end{bmatrix}$$
$$= \begin{bmatrix} (x_{1,1}w_{1,1} + \dots + x_{1,d}w_{d,1}) & \dots & (x_{1,1}w_{1,h} + \dots + x_{1,d}w_{d,h}) \\ \vdots & \ddots & \vdots \\ (x_{m,1}w_{1,1} + \dots + x_{m,d}w_{d,1}) & \dots & (x_{m,1}w_{1,h} + \dots + x_{m,d}w_{d,h}) \end{bmatrix}$$
$$= \begin{bmatrix} \sum_{i=1}^{d} x_{1,i}w_{i,1} & \dots & \sum_{i=1}^{d} x_{1,i}w_{i,h} \\ \vdots & \ddots & \vdots \\ \sum_{i=1}^{d} x_{m,i}w_{i,1} & \dots & \sum_{i=1}^{d} x_{m,i}w_{i,h} \end{bmatrix}$$



2 Matrix inverse, determinant, independence

3 Systems of linear equations

Matrix Inverse

• Definition:

A $d \times d$ square matrix **A** is called **invertible** (also **nonsingular**) if there exists a $d \times d$ square matrix **B** such that AB = BA = I (Identity matrix) given by

$$\mathbf{I} = \begin{bmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & 1 & \dots & 0 & 0 \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & \dots & 1 & 0 \\ 0 & 0 & \dots & 0 & 1 \end{bmatrix}$$

of d by d dimension

Matrix Inverse Computation

$$\mathbf{A}^{-1} = \frac{1}{\det(\mathbf{A})} \operatorname{adj}(\mathbf{A})$$

- $det(\mathbf{A})$ is the determinant of \mathbf{A}
- adj(A) is the adjugate or adjoint of A which is the transpose of its cofactor matrix C, *i.e.*, adj(A) = C[⊤]
- The cofactor $\mathbf{C}_{i,j}$ of a matrix is the (i, j)-minor $\mathbf{M}_{i,j}$ times a sign factor $(-1)^{i+j}$, *i.e.*, $\mathbf{C}_{i,j} = \mathbf{M}_{i,j} \times (-1)^{i+j}$
- $\mathbf{M}_{i,j}$ is computed through two-steps: removing the *i*-th row and *j*-th column from the original matrix to obtain a small matrix; computing the determinant of the small matrix

• For example,
$$\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$
 then $\mathbf{C} = \begin{bmatrix} d & -c \\ -b & a \end{bmatrix}$, $\operatorname{adj}(\mathbf{A}) = \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$

Determinant computation

• Example: 2×2 matrix

$$\det(\mathbf{A}) = |\mathbf{A}| = \begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc \tag{1}$$

Determinant computation

• Example: 3×3 matrix

$$|A| = \begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix} = a \begin{vmatrix} \Box & \Box & \Box \\ \Box & e & f \\ \Box & h & i \end{vmatrix} - b \begin{vmatrix} \Box & \Box & \Box \\ d & \Box & f \\ g & \Box & i \end{vmatrix} + c \begin{vmatrix} \Box & \Box & \Box \\ d & e & \Box \\ g & h & \Box \end{vmatrix}$$
$$= a \begin{vmatrix} e & f \\ h & i \end{vmatrix} - b \begin{vmatrix} d & f \\ g & i \end{vmatrix} + c \begin{vmatrix} d & e \\ g & h \end{vmatrix}$$
$$= a(ei - fh) - b(di - fg) + c(dh - eg)$$

- Each determinant of a 2 × 2 matrix in this equation is called a minor of the matrix A. This procedure can be extended to give a recursive definition for the determinant of a $n \times n$ matrix, known as **Laplace expansion**.
- Determinant has an elegant **geometric interpretation**. If interested, please refer to https://spaces.ac.cn/archives/1770.

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}.$$

$$egin{array}{c|c} a_{im} & a_{in} \ a_{jm} & a_{jn} \end{array} = \det egin{pmatrix} a_{im} & a_{in} \ a_{jm} & a_{jn} \end{pmatrix}.$$

$$\mathbf{C}=egin{pmatrix} +egin{pmatrix} a_{22}&a_{23}\ a_{32}&a_{33} \end{bmatrix} & -egin{pmatrix} a_{21}&a_{23}\ a_{31}&a_{33} \end{bmatrix} & +egin{pmatrix} a_{21}&a_{22}\ a_{31}&a_{32} \end{bmatrix} \ -egin{pmatrix} a_{12}&a_{13}\ a_{32}&a_{33} \end{bmatrix} & +egin{pmatrix} a_{11}&a_{13}\ a_{31}&a_{33} \end{bmatrix} & -egin{pmatrix} a_{11}&a_{13}\ a_{31}&a_{33} \end{bmatrix} & -egin{pmatrix} a_{11}&a_{13}\ a_{31}&a_{32} \end{bmatrix} \ +egin{pmatrix} a_{12}&a_{13}\ a_{22}&a_{23} \end{bmatrix} & -egin{pmatrix} a_{11}&a_{13}\ a_{21}&a_{23} \end{bmatrix} & +egin{pmatrix} a_{11}&a_{12}\ a_{21}&a_{22} \end{bmatrix} \end{pmatrix},$$

$$\mathbf{A} = egin{pmatrix} a_{11} & a_{12} & a_{13} \ a_{21} & a_{22} & a_{23} \ a_{31} & a_{32} & a_{33} \end{pmatrix}.$$

$$egin{array}{cc} a_{im} & a_{in} \ a_{jm} & a_{jn} \end{array} = \det egin{pmatrix} a_{im} & a_{in} \ a_{jm} & a_{jn} \end{pmatrix}.$$

$$\mathbf{C} = egin{pmatrix} + egin{pmatrix} a_{22} & a_{23} \ a_{32} & a_{33} \end{bmatrix} & -egin{pmatrix} a_{21} & a_{23} \ a_{31} & a_{33} \end{bmatrix} & +egin{pmatrix} a_{21} & a_{22} \ a_{31} & a_{33} \end{bmatrix} & -egin{pmatrix} a_{21} & a_{23} \ a_{31} & a_{33} \end{bmatrix} & -egin{pmatrix} a_{11} & a_{13} \ a_{32} & a_{33} \end{bmatrix} & -egin{pmatrix} a_{11} & a_{13} \ a_{32} & a_{33} \end{bmatrix} & -egin{pmatrix} a_{11} & a_{13} \ a_{31} & a_{33} \end{bmatrix} & -egin{pmatrix} a_{11} & a_{13} \ a_{31} & a_{33} \end{bmatrix} & -egin{pmatrix} a_{11} & a_{13} \ a_{22} & a_{23} \end{bmatrix} & -egin{pmatrix} a_{11} & a_{13} \ a_{22} & a_{23} \end{bmatrix} & +egin{pmatrix} a_{11} & a_{12} \ a_{21} & a_{22} \end{bmatrix} \end{pmatrix},$$

$$\mathbf{A} = egin{pmatrix} a_{11} & a_{12} & a_{13} \ a_{21} & a_{22} & a_{23} \ a_{31} & a_{32} & a_{33} \end{pmatrix}. \qquad \qquad \begin{vmatrix} a_{im} & a_{in} \ a_{jm} & a_{jn} \end{vmatrix} = \det egin{pmatrix} a_{im} \ a_{jm} \ a_{jm} \end{vmatrix}$$

 a_{in}

 a_{jr}

$$\mathbf{C}=egin{pmatrix} +egin{pmatrix} a_{22}&a_{23}\ a_{32}&a_{33}\end{bmatrix} & -egin{pmatrix} a_{21}&a_{23}\ a_{31}&a_{33}\end{bmatrix} & +egin{pmatrix} a_{21}&a_{22}\ a_{31}&a_{32}\end{bmatrix} \ -egin{pmatrix} a_{12}&a_{13}\ a_{32}&a_{33}\end{bmatrix} & +egin{pmatrix} a_{11}&a_{13}\ a_{31}&a_{33}\end{bmatrix} & -egin{pmatrix} a_{11}&a_{13}\ a_{31}&a_{33}\end{bmatrix} & -egin{pmatrix} a_{11}&a_{13}\ a_{31}&a_{32}\end{bmatrix} \ +egin{pmatrix} a_{12}&a_{13}\ a_{22}&a_{23}\end{bmatrix} & -egin{pmatrix} a_{11}&a_{13}\ a_{21}&a_{23}\end{bmatrix} & +egin{pmatrix} a_{11}&a_{12}\ a_{21}&a_{22}\end{bmatrix} \end{pmatrix},$$

$$\mathbf{A} = egin{pmatrix} a_{11} & a_{12} & a_{13} \ a_{21} & a_{22} & a_{23} \ a_{31} & a_{32} & a_{33} \end{pmatrix}. \qquad \qquad \begin{vmatrix} a_{im} & a_{in} \ a_{jm} & a_{jn} \end{vmatrix} = \det egin{pmatrix} a_{im} & a_{in} \ a_{jm} & a_{jn} \end{vmatrix}$$

).

$$\mathbf{C} = egin{pmatrix} + egin{pmatrix} a_{22} & a_{23} \ a_{32} & a_{33} \ \end{pmatrix} egin{pmatrix} - egin{pmatrix} a_{21} & a_{22} \ a_{31} & a_{33} \ \end{pmatrix} egin{pmatrix} + egin{pmatrix} a_{21} & a_{22} \ a_{31} & a_{32} \ \end{pmatrix} egin{pmatrix} - egin{pmatrix} a_{12} & a_{13} \ a_{32} \ a_{33} \ \end{pmatrix} egin{pmatrix} + egin{pmatrix} a_{21} & a_{22} \ a_{31} & a_{32} \ \end{pmatrix} egin{pmatrix} + egin{pmatrix} a_{21} & a_{23} \ a_{31} & a_{33} \ \end{pmatrix} egin{pmatrix} + egin{pmatrix} a_{21} & a_{22} \ a_{31} & a_{32} \ \end{pmatrix} egin{pmatrix} + egin{pmatrix} a_{21} & a_{23} \ a_{31} & a_{32} \ \end{pmatrix} egin{pmatrix} + egin{pmatrix} a_{21} & a_{23} \ a_{31} & a_{32} \ \end{pmatrix} egin{pmatrix} + egin{pmatrix} a_{21} & a_{22} \ a_{23} \ \end{pmatrix} egin{pmatrix} + egin{pmatrix} a_{21} & a_{23} \ a_{21} & a_{22} \ \end{pmatrix} egin{pmatrix} + egin{pmatrix} a_{21} & a_{22} \ a_{21} & a_{22} \ \end{pmatrix} egin{pmatrix} + egin{pmatrix} a_{21} & a_{22} \ a_{21} & a_{22} \ \end{pmatrix} egin{pmatrix} + egin{pmatrix} a_{21} & a_{22} \ a_{21} & a_{22} \ \end{pmatrix} egin{pmatrix} + egin{pmatrix} a_{21} & a_{22} \ a_{23} \ \end{pmatrix} egin{pmatrix} + egin{pmatrix} a_{21} & a_{22} \ a_{21} & a_{22} \ \end{pmatrix} egin{pmatrix} + egin{pmatrix} a_{21} & a_{22} \ a_{21} & a_{22} \ \end{pmatrix} egin{pmatrix} + egin{pmatrix} a_{21} & a_{22} \ a_{21} & a_{22} \ \end{pmatrix} egin{pmatrix} + egin{pmatrix} a_{21} & a_{22} \ a_{21} & a_{22} \ \end{pmatrix} egin{pmatrix} + egin{pmatrix} a_{21} & a_{22} \ a_{21} & a_{22} \ \end{pmatrix} egin{pmatrix} + egin{pmatrix} a_{21} & a_{22} \ a_{21} & a_{22} \ \end{pmatrix} egin{pmatrix} + egin{pmatrix} a_{21} & a_{22} \ a_{21} & a_{22} \ \end{pmatrix} egin{pmatrix} + egin{pmatrix} a_{21} & a_{22} \ a_{21} & a_{22} \ \end{pmatrix} egin{pmatrix} + egin{pmatrix} a_{21} & a_{22} \ a_{21} & a_{22} \ \end{pmatrix} egin{pmatrix} + egin{pmatrix} a_{21} & a_{22} \ a_{21} & a_{22} \ \end{pmatrix} egin{pmatrix} + egin{pmatrix} a_{21} & a_{22} \ a_{21} & a_{22} \ \end{pmatrix} egin{pmatrix} + egin{pmatrix} a_{21} & a_{22} \ a_{21} & a_{22} \ \end{pmatrix} egin{pmatrix} + egin{pmatrix} a_{21} & a_{22} \ a_{21} & a_{22} \ \end{pmatrix} egin{pmatrix} + egin{pmatrix} a_{21} & a_{22} \ a_{21} & a_{22} \ \end{pmatrix} egin{pmatrix} a_{21} & a$$

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}. \qquad \qquad \begin{vmatrix} a_{im} & a_{in} \\ a_{jm} & a_{jn} \end{vmatrix} = \det \begin{pmatrix} a_{im} & a_{in} \\ a_{jm} & a_{jn} \end{pmatrix}$$

$$\mathbf{C}=egin{pmatrix} +egin{pmatrix} a_{22}&a_{23}\ a_{32}&a_{33}\end{bmatrix} & -egin{pmatrix} a_{21}&a_{23}\ a_{31}&a_{33}\end{bmatrix} & +egin{pmatrix} a_{21}&a_{22}\ a_{31}&a_{32}\end{bmatrix} \ -egin{pmatrix} a_{12}&a_{13}\ a_{32}&a_{33}\end{bmatrix} & +egin{pmatrix} a_{11}&a_{13}\ a_{31}&a_{33}\end{bmatrix} & -egin{pmatrix} a_{11}&a_{13}\ a_{31}&a_{33}\end{bmatrix} & -egin{pmatrix} a_{11}&a_{13}\ a_{31}&a_{32}\end{bmatrix} \ +egin{pmatrix} a_{12}&a_{13}\ a_{22}&a_{23}\end{bmatrix} & -egin{pmatrix} a_{11}&a_{13}\ a_{21}&a_{23}\end{bmatrix} & +egin{pmatrix} a_{11}&a_{12}\ a_{21}&a_{22}\end{bmatrix} \end{pmatrix},$$

Linear dependence and independence

• A collection of *d*-vectors $\mathbf{x}_1, \ldots, \mathbf{x}_m$ (with $m \ge 1$) is called linearly dependent if

$$\beta_1 \mathbf{x}_1 + \ldots + \beta_m \mathbf{x}_m = 0$$

holds for some β_1, \ldots, β_m that are not all zero.

• A collection of *d*-vectors $\mathbf{x}_1, \ldots, \mathbf{x}_m$ (with $m \ge 1$) is called linearly independent, which means that

$$\beta_1 \mathbf{x}_1 + \ldots + \beta_m \mathbf{x}_m = 0$$

only holds for $\beta_1 = \ldots = \beta_m = 0$.

Linear dependence and independence

Geometry of dependency and independency





2 Matrix inverse, determinant, independence



• Consider a system of m linear equations in d variables w_1, \ldots, w_d :

 $x_{1,1}w_1 + x_{1,2}w_2 + \dots + x_{1,d}w_d = y_1$

 $x_{2,1}w_1 + x_{2,2}w_2 + \dots + x_{2,d}w_d = y_2$

:

 $x_{m,1}w_1 + x_{m,2}w_2 + \ldots + x_{m,d}w_d = y_m$

These equations can be written compactly in matrix-vector notation:

$$\mathbf{X}\mathbf{w} = \mathbf{y},$$

where

$$\mathbf{X} = \begin{bmatrix} x_{1,1} & \dots & x_{1,d} \\ \vdots & \ddots & \vdots \\ x_{m,1} & \dots & x_{m,d} \end{bmatrix}, \qquad \mathbf{w} = \begin{bmatrix} w_1 \\ \vdots \\ w_d \end{bmatrix}, \qquad \mathbf{y} = \begin{bmatrix} y_1 \\ \vdots \\ y_m \end{bmatrix}$$

Note: **X** is of $m \times d$ dimension.

(i) Square of even-determined system: m = d in $\mathbf{X}\mathbf{w} = \mathbf{y}, \mathbf{X} \in \mathbb{R}^{m \times d}$ (equal number of equations and unknowns, *i.e.*, $\mathbf{X} \in \mathbb{R}^{d \times d}$)

If **X** is invertible (or $\mathbf{X}^{-1}\mathbf{X} = \mathbf{I}$), then pre-multiply both sides by \mathbf{X}^{-1} , we have

 $\mathbf{X}^{-1}\mathbf{X}\mathbf{w} = \mathbf{X}^{-1}\mathbf{y}$ $\Rightarrow \mathbf{w} = \mathbf{X}^{-1}\mathbf{y}$

If all rows or columns of **X** are **linearly independent**, then **X** is **invertible**.

Systems of linear equations

Example $w_1 + w_2 = 4$ (1) Two unknowns and $w_1 - 2w_2 = 1$ (2) two equations Equation (1) $\begin{bmatrix} 1 & 1 \\ 1 & -2 \end{bmatrix} \qquad \begin{bmatrix} w_1 \\ w_2 \end{bmatrix} = \begin{bmatrix} 4 \\ 1 \end{bmatrix} \stackrel{25}{}_2$ X w Equation (2) w_2 0.5 $\mathbf{w} = \mathbf{X}^{-1}\mathbf{v}$ -0.5 $= \begin{bmatrix} 1 & 1 \\ 1 & -2 \end{bmatrix}^{-1} \begin{bmatrix} 4 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$ -1 -1.5 -2 -3 w_1

Here, the rows or columns of **X** are **linearly independent**, hence **X** is **invert-ible**.

(ii)Over-determined system: m > d in $\mathbf{X}\mathbf{w} = \mathbf{y}, \mathbf{X} \in \mathbb{R}^{m \times d}$ (i.e., there are more equations than unknowns)

- This set of linear equations has NO exact solution (X is non-square and hence not invertible). However, an **approximated solution** is yet available.
- If the left-inverse of X exists such that $X^{\dagger}X = I$, then pre-multiply both sides by X^{\dagger} results in

 $\begin{aligned} \mathbf{X}^{\dagger} \mathbf{X} \mathbf{w} &= \mathbf{X}^{\dagger} \mathbf{y} \\ \Rightarrow \mathbf{w} &= \mathbf{X}^{\dagger} \mathbf{y} \end{aligned}$

- Definition: a matrix **B** that satisfies BA=I (identity matrix) is called a left-inverse of **A**. (Note: **A** is m-by-d and **B** is d-by-m.
- Note: The left-inverse can be computed as $\mathbf{X}^{\dagger} = (\mathbf{X}^{\top}\mathbf{X})^{-1}\mathbf{X}^{\top}$ given $\mathbf{X}^{\top}\mathbf{X}$ is invertible.

Systems of linear equations

Example
$$w_1 + w_2 = 1$$
 (1)
 $w_1 - w_2 = 0$ (2)
 $w_1 = 2$ (3)

Two unknowns and three equations

$$\begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} w_1 \\ w_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}$$

X w y

This set of linear equations has NO exact solution.



 $\mathbf{w} = \mathbf{X}^{\dagger} \mathbf{y} = (\mathbf{X}^{\top} \mathbf{X})^{-1} \mathbf{X}^{\top} \mathbf{y} \quad \text{Here } \mathbf{X}^{\top} \mathbf{X} \text{ is invertible.}$

 $= \begin{bmatrix} 3 & 0 \\ 0 & 2 \end{bmatrix}^{-1} \begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 0.5 \end{bmatrix}$ (Approximation)

(iii)Under-determined system: m < d in $\mathbf{X}\mathbf{w} = \mathbf{y}, \mathbf{X} \in \mathbb{R}^{m \times d}$ (i.e., there are more unknowns than equations \Rightarrow infinite number of solutions)

If the right-inverse of X exists such that XX[†] = I, then the *d*-vector w = X[†]y (one of the infinite cases) satisfies the equation Xw = y, *i.e.*,

$$\mathbf{X}\mathbf{X}^{\dagger}\mathbf{y} = \mathbf{y}$$

$$\Rightarrow \mathbf{y} = \mathbf{y}$$

- **Definition**: a matrix **B** that satisfies AB=I (identity matrix) is called a right-inverse of **A**. (Note: **A** is m-by-d and **B** is d-by-m).
- Note: The right-inverse can be computed as $\mathbf{X}^{\dagger} = \mathbf{X}^{\top} (\mathbf{X} \mathbf{X}^{\top})^{-1}$ given $\mathbf{X} \mathbf{X}^{\top}$ is invertible.

Derivation:

- Under-determined system: m < d in Xw = y, X ∈ ℝ^{m×d}
 (i.e., there are more unknowns than equations ⇒ infinite number of solutions ⇒ but a unique solution is yet possible by constraining the search using w = X^Ta !)
- If $\mathbf{X}\mathbf{X}^{\top}$ is invertible, let $\mathbf{w} = \mathbf{X}^{\top}\mathbf{a}$, then

$$\mathbf{X}\mathbf{X}^{\top}\mathbf{a} = \mathbf{y}$$

$$\Rightarrow \mathbf{a} = (\mathbf{X}\mathbf{X}^{\top})^{-1}\mathbf{y}$$

$$\mathbf{w} = \mathbf{X}^{\top}\mathbf{a} = \underbrace{\mathbf{X}^{\top}(\mathbf{X}\mathbf{X}^{\top})^{-1}}_{\mathbf{X}^{\dagger}}\mathbf{y}$$

Systems of linear equations

Example
$$w_1 + 2w_2 + 3w_3 = 2$$
 (1) Three unknowns and
 $w_1 - 2w_2 + 3w_3 = 1$ (2) two equations

$$\begin{bmatrix} 1 & 2 & 3 \\ 1 & -2 & 3 \end{bmatrix} \begin{bmatrix} w_1 \\ w_2 \\ w_3 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

$$X \qquad W \qquad Y$$

This set of linear equations has infinitely many solutions along the intersection line.



36 / 37

 $\mathbf{w} = \mathbf{X}^{\top} (\mathbf{X} \mathbf{X}^{\top})^{-1} \mathbf{y} \quad \text{Here } \mathbf{X} \mathbf{X}^{\top} \text{ is invertible.}$ $= \begin{bmatrix} 1 & 1 \\ 2 & -2 \\ 3 & 3 \end{bmatrix} \begin{bmatrix} 14 & 6 \\ 6 & 14 \end{bmatrix}^{-1} \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 0.15 \\ 0.25 \\ 0.45 \end{bmatrix} \quad \text{(Constrained solution)}$

